

Characterisations of Intersection Graphs by Vertex Orderings^{*}

David R. Wood

School of Computer Science
Carleton University, Ottawa, Canada

Department of Applied Mathematics
Charles University, Prague, Czech Republic

davidw@scs.carleton.ca

February 1, 2008

Abstract

Characterisations of interval graphs, comparability graphs, co-comparability graphs, permutation graphs, and split graphs in terms of linear orderings of the vertex set are presented. As an application, it is proved that interval graphs, co-comparability graphs, AT-free graphs, and split graphs have bandwidth bounded by their maximum degree.

1 Introduction

We consider finite, simple and undirected graphs G with vertex set $V(G)$, edge set $E(G)$, and maximum degree $\Delta(G)$. The *complement* of G is the graph $\overline{G}$ with vertex set $V(G)$ and edge set $\{vw : v, w \in V(G), vw \notin E(G)\}$. A *vertex ordering* of G is a total order $(v_1, v_2, \dots, v_n)$ of $V(G)$. Let $\mathcal{S}$ be a finite family of sets. The *intersection graph* of $\mathcal{S}$ has vertex set $\mathcal{S}$ and edge set $\{AB : A, B \in \mathcal{S}, A \cap B \neq \emptyset\}$. This paper presents characterisations of a number of popular intersection graphs in terms of vertex orderings.

In a vertex ordering $(v_1, v_2, \dots, v_n)$ of a graph G , the *width* of an edge $v_i v_j \in E(G)$ is $|i - j|$. The maximum width of an edge is the *width* of the ordering. The *bandwidth* of G is the minimum width of a vertex ordering of G . Bandwidth is a ubiquitous concept with numerous applications (see [2]). Obviously the bandwidth of G is at least $\frac{1}{2}\Delta(G)$. As an application of our results, we prove upper bounds on the bandwidth of many intersection graphs G in terms of $\Delta(G)$.

^{*}Research supported by NSERC and COMBSTRU.

2 Interval Graphs

An *interval graph* is the intersection graph of a finite set of closed intervals in $\mathbb{R}$. We have the following characterisation of interval graphs.

Theorem 1. *A graph G is an interval graph if and only if G has a vertex ordering $(v_1, v_2, \dots, v_n)$ such that*

$$\forall i < j < k, v_i v_k \in E(G) \Rightarrow v_i v_j \in E(G) . \quad (1)$$

Proof. Let $(v_1, v_2, \dots, v_n)$ be a vertex ordering of G satisfying (1). For each vertex v_i , associate the interval $[i, r(i)]$, where $v_{r(i)}$ is the rightmost neighbour of v_i . For every edge $v_i v_j \in E(G)$ with $i < j$, $j \in [i, r(i)] \cap [j, r(j)]$. For every non-edge $v_i v_j \in E(\overline{G})$ with $i < j$, we have $r(i) < j$ by (1), and $[i, r(i)] \cap [j, r(j)] = \emptyset$. Hence the intersection graph of $\{[i, r(i)] : v_i \in V(G)\}$ is G .

Let G be an interval graph. It is well known that we can assume that the endpoints of the intervals are distinct. Let $(v_1, v_2, \dots, v_n)$ be a vertex ordering of G determined by increasing values of the left endpoints of the intervals. For all $i < j < k$, the left endpoint of v_j is between the left endpoints of v_i and v_k . If $v_i v_k \in E(G)$ then the right endpoint of v_i is to the right of the left endpoint of v_k . Thus the left endpoint of v_j is in the interval for v_i . Hence $v_i v_j \in E(G)$, as claimed. $\square$

As far as we are aware, Theorem 1 has not appeared in the literature, although similar results are known. For example, Gilmore and Hoffman [10] proved that G is an interval graph if and only if there is an ordering of the maximal cliques of G such that for each vertex v , the maximal cliques containing v appear consecutively.

Theorem 1 implies the following result of Fomin and Golovach [8].

Corollary 1 ([8]). *Every interval graph G has bandwidth at most $\Delta(G)$.*

Proof. In the vertex ordering $(v_1, v_2, \dots, v_n)$ from Theorem 1, the width of an edge $v_i v_k \in E(G)$ is $|\{v_i v_j \in E(G) : i < j \leq k\}| \leq \deg(v_i) \leq \Delta(G)$. $\square$

A *proper interval graph* is the intersection graph of a finite set $\mathcal{S}$ of closed intervals in $\mathbb{R}$ such that $A \not\subset B$ for all $A, B \in \mathcal{S}$. The following characterisation is due to Looges and Olariu [15] (see [3]).

Theorem 2 ([15]). *A graph G is a proper interval graph if and only if G has a vertex ordering $(v_1, v_2, \dots, v_n)$ such that,*

$$\forall i < j < k, v_i v_k \in E(G) \Rightarrow v_i v_j \in E(G) \wedge v_j v_k \in E(G) . \quad (2)$$

It is easily seen that the bandwidth of a proper interval graph is one less than the maximum clique size. Moreover, Kaplan and Shamir [12] proved that the bandwidth of any graph G equals the minimum, taken over all proper interval supergraphs G' of G , of the bandwidth of G' .

3 Comparability Graphs

Let $(P, \preceq)$ be a poset. The *comparability graph* of $(P, \preceq)$ has vertex set P , and distinct elements are adjacent if and only if they are comparable under $\preceq$. We have the following characterisation of comparability graphs.

Theorem 3. *The following are equivalent for a graph G :*

- (a) G is a comparability graph,
- (b) G has a vertex ordering $(v_1, v_2, \dots, v_n)$ such that,

$$\forall i < j < k, v_i v_j \in E(G) \wedge v_j v_k \in E(G) \Rightarrow v_i v_k \in E(G) . \quad (3)$$

Proof. Let G be the comparability graph of a poset $(V(G), \preceq)$. A linear extension of $\preceq$ satisfies (3). Given a vertex ordering that satisfies (3), define $v_i \prec v_j$ whenever $v_i v_j \in E(G)$ and $i < j$. Thus $(V(G), \preceq)$ is a poset, and G is a comparability graph. $\square$

A *co-comparability graph* is a complement of a comparability graph. As illustrated in Figure 1, a *function diagram* is a set $\{c_i : 1 \leq i \leq n\}$, where for each c_i is a curve $\{(x, f_i(x)) : 0 \leq x \leq 1\}$ for some function $f_i : [0, 1] \rightarrow \mathbb{R}$. If each c_i is a line segment we say $\{c_i : 1 \leq i \leq n\}$ is *linear*.




Figure 1: A vertex ordering of the intersection graph of a function diagram.

Theorem 4. *The following are equivalent for a graph G :*

- (a) G is a co-comparability graph,
- (b) G is the intersection graph of a function diagram, and
- (c) G has a vertex ordering $(v_1, v_2, \dots, v_n)$ such that,

$$\forall i < j < k, v_i v_k \in E(G) \Rightarrow v_i v_j \in E(G) \vee v_j v_k \in E(G) . \quad (4)$$

Proof. Kratochvíl *et al.* [14] and Golumbic *et al.* [11] independently proved that (a) and (b) are equivalent.

We now prove that (c) implies (a). Suppose that G has a vertex ordering $(v_1, v_2, \dots, v_n)$ satisfying (4). Define $v_i \prec v_j$ if $i < j$ and $v_i v_j \notin E(G)$. Obviously $\prec$ is antisymmetric. Suppose $v_i \prec v_j$ and $v_j \prec v_k$. Then $i < k$ and $v_i v_k \notin E(G)$, as otherwise (4) fails. That

is, $v_i \prec v_k$. Thus $\prec$ is transitive, and $(V(G), \preceq)$ is a poset, whose comparability graph is $\overline{G}$. Therefore G is a co-comparability graph.

We now prove that (b) implies (c). Let G be the intersection graph of a function diagram $\{c_i : 1 \leq i \leq n\}$ with corresponding functions $\{f_i : 1 \leq i \leq n\}$. Re-index so that $f_i(0) \leq f_{i+1}(0)$ for all $1 \leq i \leq n-1$. Associate a vertex v_i with each function f_i . Consider an edge $v_i v_k \in E(G)$ and a vertex v_j with $i < j < k$. There is a region S bounded by c_i , c_k , and the line $X = 0$, such that c_j intersects the closed interior of S and the closed exterior of S . Thus c_j intersects the boundary of S . Since f_j is a function on $[0, 1]$, c_j intersects the boundary of S at a point on c_i or c_k . Thus $c_i \cap c_j \neq \emptyset$ or $c_j \cap c_k \neq \emptyset$. Hence $v_i v_k \in E(G)$ or $v_j v_k \in E(G)$. That is, the vertex ordering $(v_1, v_2, \dots, v_n)$ satisfies (4). Note that we could have ordered the vertices with respect to any fixed value of $x_0 \in [0, 1]$, and in general, there are many vertex orderings that satisfy (4). $\square$

Corollary 2. *Every co-comparability graph G has bandwidth at most $2\Delta(G) - 1$.*

Proof. In the vertex ordering $(v_1, v_2, \dots, v_n)$ from Theorem 4, the width of an edge $v_i v_k \in E(G)$ is at most $|\{v_i v_j \in E(G) : i < j < k\}| + |\{v_j v_k \in E(G) : i < j < k\}| + 1 \leq (\deg(v_i) - 1) + (\deg(v_k) - 1) + 1 \leq 2\Delta(G) - 1$. $\square$

It is interesting to ask whether Corollary 2 is tight. It is easily seen that the complete bipartite graph $K_{n,n}$, which is a co-comparability graph with maximum degree n , has bandwidth $3n/2$.

Let π be a permutation of $\{1, 2, \dots, n\}$. Let $\pi^{-1}(i)$ denote the position of i in π . The *permutation graph* associated with π has vertex set $\{v_1, v_2, \dots, v_n\}$ and edge set $\{v_i v_j : (i - j)(\pi^{-1}(i) - \pi^{-1}(j)) < 0\}$. The following characterisations of permutation graphs can be derived from results of Dushnik and Miller [6] and Baker *et al.* [1]. Part (e) is proved as in Theorems 3 and 4.

Theorem 5 ([1, 6]). *The following are equivalent for a graph G :*

- (a) G is a permutation graph,
- (b) G is the intersection graph of a linear function diagram,
- (c) G is a comparability graph and a co-comparability graph,
- (d) G is the comparability graph of a two-dimensional poset,
- (e) G has a vertex ordering that simultaneously satisfies (3) and (4).

4 AT-free Graphs

An *asteroidal triple* in a graph consists of an independent set of three vertices such that each pair is joined by a path that avoids the neighbourhood of the third. A graph is *asteroidal triple-free* (*AT-free*) if it contains no asteroidal triples.

Lemma 1. *Every AT-free graph G has bandwidth at most $3\Delta(G)$.*

Proof. A *caterpillar* is a tree for which a path (called the spine) is obtained by deleting all the leaves. Let $(v_1, v_2, \dots, v_m)$ be the spine of a caterpillar T . The vertex ordering of T obtained by inserting the leaves adjacent to each v_i immediately after v_i has bandwidth at most $\Delta(T)$.

Kloks *et al.* [13] proved that every (connected) AT-free graph G has a spanning caterpillar subgraph T , and adjacent vertices in G are at distance at most four in T . Moreover, for any edge $vw \in E(G)$ with v and w at distance four in T , both v and w are leaves of T . Consider the above vertex ordering of T to be a vertex ordering of G . The bandwidth is at most $3\Delta(T) \leq 3\Delta(G)$. $\square$

5 Chordal Graphs

A *chord* of a cycle C is an edge not in C connecting two vertices in C . A graph is *chordal* if every induced cycle on at least four vertices has at least one chord. The following famous characterisation of chordal graphs is due to Dirac [5], Fulkerson and Gross [9], and Rose [16].

Theorem 6 ([5, 9, 16]). *The following are equivalent for a graph G :*

- (a) G is a chordal,
- (b) G is the intersection graph of subtrees of a tree, and
- (c) G has a vertex ordering $(v_1, v_2, \dots, v_n)$ such that,

$$\forall i < j < k, v_i v_j \in E(G) \wedge v_i v_k \in E(G) \Rightarrow v_j v_k \in E(G) . \quad (5)$$

A striking generalisation of Theorem 6 for k -chordal graphs is given by Dendris *et al.* [4]. A vertex ordering satisfying (5) is called a *perfect elimination* vertex ordering. It is not possible to bound the bandwidth of every chordal graph G in terms of $\Delta(G)$. For example, the bandwidth of the complete binary tree on n vertices is $\approx n/\log n$ [17].

A graph G is a *split graph* if $V(G) = K \cup I$, where K induces a complete graph of G , and I is an independent set of G .

Theorem 7. *The following are equivalent for a graph G :*

- (a) G is a split graph,
- (b) G is chordal and $\overline{G}$ is chordal,
- (c) G has a vertex ordering $(v_1, v_2, \dots, v_n)$ simultaneously satisfying (5) and

$$\forall i < j < k, v_i v_j \in E(G) \Rightarrow v_j v_k \in E(G) \vee v_i v_k \in E(G) , \quad (6)$$

- (d) G has a vertex ordering $(v_1, v_2, \dots, v_n)$ such that,

$$\forall i < j < k, v_i v_j \in E(G) \Rightarrow v_j v_k \in E(G) . \quad (7)$$

Proof. Földes and Hammer [7] proved that (a) and (b) are equivalent.

Observe that (d) implies (c) trivially. We now prove that (a) implies (d). Let G be a split graph with $V(G) = K \cup I$, where K induces a complete subgraph and I is an

independent set. Let $m = |I|$. Consider a vertex ordering $(v_1, v_2, \dots, v_n)$ of G where $I = \{v_1, v_2, \dots, v_m\}$ and $K = \{v_{m+1}, v_{m+2}, \dots, v_n\}$. Suppose that $1 \leq i < j < k \leq n$ and $v_i v_j \in E(G)$. There is no edge with both endpoints in I . Thus $j \geq m + 1$, and both $v_j, v_k \in K$. Hence $v_j v_k \in E(G)$, and $(v_1, v_2, \dots, v_n)$ satisfies (7).

It remains to prove that (c) implies (b). Let $(v_1, v_2, \dots, v_n)$ be a vertex ordering of a graph G satisfying (5) and (6). By Theorem 6, G is chordal. Equation (6) is equivalent to:

$$\forall i < j < k, v_j v_k \in E(\overline{G}) \wedge v_i v_k \in E(\overline{G}) \Rightarrow v_i v_j \in E(\overline{G}) . \quad (8)$$

That is, $(v_n, v_{n-1}, \dots, v_1)$ is a perfect elimination vertex ordering of $\overline{G}$. By Theorem 6, $\overline{G}$ is chordal. $\square$

Lemma 2. *Every split graph G has bandwidth at most $\Delta(G) (\Delta(G) + 2)$. For all $\Delta \geq 2$ there is a split graph G with $\Delta(G) = \Delta$, and G has bandwidth at least $\Delta(G)^2/12$.*

Proof. First we prove the upper bound. Let G be a split graph with $V(G) = K \cup I$, where K induces a complete subgraph, and I is an independent set. The result is trivial if $K = \emptyset$. Now assume that $K \neq \emptyset$. Let I_0 be the set of isolated vertices in G . Consider a vertex ordering π in which the vertices in I_0 precede all other vertices. Let $I_1 = I \setminus I_0$. Regardless of the order of $I_1 \cup K$, the bandwidth of π is at most $|I_1| + |K| - 1$. Thus it suffices to prove that $|I_1| + |K| \leq \Delta(G) (\Delta(G) + 2) + 1$.

If $I_1 = \emptyset$ then π has bandwidth $\Delta(G)$. Now assume that $I_1 \neq \emptyset$. Let a be the average degree of vertices in I_1 . Thus $1 \leq a \leq |K|$. For each vertex $v \in K$, let $b_v = \deg(v) - |K| + 1$. That is, b_v is the number of edges between v and I_1 . Let $b = \sum_{v \in K} b_v / |K|$. Thus $a|I_1| = b|K|$, which implies that

$$|I_1| + |K| = \frac{b|K|}{a} + |K| = \frac{(b+a)|K|}{a} .$$

Now $\Delta(G)$ is at least the average degree of the vertices in K . That is, $\Delta(G) \geq |K| - 1 + b$. Hence

$$\frac{|I_1| + |K|}{(\Delta(G) + 1)^2} \leq \frac{(b+a)|K|}{a(|K| + b)^2} .$$

Since $a \leq |K|$,

$$\frac{|I_1| + |K|}{(\Delta(G) + 1)^2} \leq \frac{|K|}{a(|K| + b)} .$$

Since $a \geq 1$ and $b \geq 0$,

$$|I_1| + |K| \leq (\Delta(G) + 1)^2 = \Delta(G) (\Delta(G) + 2) + 1 ,$$

as required.

Now we prove the lower bound. Given Δ , let $n = \lfloor \Delta/2 \rfloor$. Let G be the split graph with $V(G) = K \cup I$, where K is a complete graph on n vertices, and I is an independent set on $n(\Delta - n + 1)$ vertices, such that every vertex in K is adjacent to $\Delta - n + 1$ vertices in I , and every vertex in I is adjacent to exactly one vertex in K . Clearly G has diameter

3, maximum degree Δ , and $n + n(\Delta - n + 1) = n(\Delta + n - 2)$ vertices. It is easily seen that every connected graph with n' vertices and diameter d' has bandwidth at least $(n' - 1)/d'$ [2]. Thus G has bandwidth at least

$$\frac{1}{3}(n(\Delta - n + 2) - 1) = \frac{1}{3}\left(\left\lfloor \frac{\Delta}{2} \right\rfloor \left\lceil \frac{\Delta + 2}{2} \right\rceil - 1\right) \geq \frac{\Delta^2}{12}.$$

□

References

- [1] K. A. BAKER, P. C. FISHBURN, AND F. S. ROBERTS, Partial orders of dimension 2. *Networks*, **2**:11–28, 1972.
- [2] P. Z. CHINN, J. CHVÁTALOVÁ, A. K. DEWDNEY, AND N. E. GIBBS, The bandwidth problem for graphs and matrices—a survey. *J. Graph Theory*, **6**(3):223–254, 1982.
- [3] D. G. CORNEIL, A simple 3-sweep LBFS algorithm for the recognition of unit interval graphs. *Discrete Applied Math.*, **139**:371–379, 2004.
- [4] N. D. DENDRIS, L. M. KIROUSIS, AND D. M. THILIKOS, Fugitive-search games on graphs and related parameters. *Theoret. Comput. Sci.*, **172**(1-2):233–254, 1997.
- [5] G. A. DIRAC, On rigid circuit graphs. *Abh. Math. Sem. Univ. Hamburg*, **25**:71–76, 1961.
- [6] B. DUSHNIK AND E. W. MILLER, Partially ordered sets. *Amer. J. Math.*, **63**:600–610, 1941.
- [7] S. FÖLDES AND P. L. HAMMER, Split graphs. In *Proc. 8th Southeastern Conference on Combinatorics, Graph Theory and Computing*, vol. 19 of *Congressus Numerantium*, pp. 311–315, Utilitas Math., 1977.
- [8] F. V. FOMIN AND P. A. GOLOVACH, Interval degree and bandwidth of a graph. *Discrete Appl. Math.*, **129**(2-3):345–359, 2003.
- [9] D. R. FULKERSON AND O. A. GROSS, Incidence matrices and interval graphs. *Pacific J. Math.*, **15**:835–855, 1965.
- [10] P. C. GILMORE AND A. J. HOFFMAN, A characterization of comparability graphs and of interval graphs. *Canad. J. Math.*, **16**:539–548, 1964.
- [11] M. C. GOLUMBIC, D. ROTEM, AND J. URRUTIA, Comparability graphs and intersection graphs. *Discrete Math.*, **43**:37–46, 1983.

- [12] H. KAPLAN AND R. SHAMIR, Pathwidth, bandwidth, and completion problems to proper interval graphs with small cliques. *SIAM J. Comput.*, **25(3)**:540–561, 1996.
- [13] T. KLOKS, D. KRATSCH, AND H. MÜLLER, Approximating the bandwidth for asteroidal triple-free graphs. *J. Algorithms*, **32(1)**:41–57, 1999.
- [14] J. KRATOCHVÍL, M. GOLJAN, AND P. KUČERA, String graphs. *Rozprawy Československé Akad. Věd Řada Mat. Přírod. Věd*, **96(3)**, 1986.
- [15] P. J. LOOGES AND S. OLARIU, Optimal greedy algorithms for indifference graphs. *Comput. Math. Appl.*, **25(7)**:15–25, 1993.
- [16] D. J. ROSE, Triangulated graphs and the elimination process. *J. Math. Anal. Appl.*, **32**:597–609, 1970.
- [17] L. SMITHLINE, Bandwidth of the complete k -ary tree. *Discrete Math.*, **142(1-3)**:203–212, 1995.